

Effective Potential for the Second Virial Coefficient at Feshbach Resonance

Rajat K. Bhaduri¹ and M. K. Srivastava²

1. Department of Physics and Astronomy, McMaster University, Hamilton, Canada L8S 4M1

2. Institute Instrumentation Centre, I.I.T., Roorkee 247667, India

Abstract

In a recent paper entitled “High temperature expansion applied to fermions near Feshbach resonance”, (Phys. Rev. Lett. **92** 160404 (2004)), Ho and Mueller have demonstrated a remarkable similarity between its high and low temperature properties at resonance. The quantum second virial coefficient plays a crucial role in their analysis, and has a universal value at resonance. In this paper, we explore the connection between the quantum and classical second virial coefficients, and show that near a Feshbach resonance an exact mapping from the quantum to classical form is possible. This gives rise to a scale-independent inverse square effective potential for the classical virial coefficient. It is suggested that this may be tested by measuring the isothermal compressibility of the gas on the repulsive side of the resonance..

PACS: 03.75.-b, 03.75.Ss

Great advances have been made in the study of ultracold trapped atoms in recent times. The interaction between the atoms is normally weak in a dilute gas, but may be enhanced drastically by making use of the so-called Feshbach resonance [1]. This may be achieved by applying a magnetic field to tune the energy of the Zeeman level of the trapped atoms to one of the molecular resonances. At a zero-energy resonance, the scattering phase shift is $\pi/2$, and the cross section reaches the unitary limit. It is argued that the properties of a gas at Feshbach resonance have a universal character [2], irrespective of the type of gas in the trap. In this connection, Ho and Mueller [3] recently considered the interaction energy of a two-component Fermi gas near a Feshbach resonance. They compared the available experimental data of the ultracold gas in the vicinity of a Feshbach resonance with the predictions in the Boltzmann regime using the *quantum* second virial coefficient. Surprisingly, it was found that the high temperature expression, when used at micro-Kelvin temperatures, explains the data reasonably well. In the present work, we examine the use of the quantal second virial coefficient in more detail, and its connection to its classical counterpart at the Feshbach resonance. We find that the quantal expression may be mapped onto the classical formula exactly in the vicinity of the Feshbach resonance using an inverse square interaction in the configuration space.

To appreciate this in more detail, note that the virial expansion of a (one-component) gas (classical as well as quantal) for the pressure P at a temperature T is given by [4]

$$\frac{P}{n\tau} = 1 + a_2(n\lambda^3) + a_3(n\lambda^3)^2 + \dots, \quad (1)$$

where $a_2, a_3, \dots$ etc are the dimensionless second, third..virial coefficients, $\tau = k_B T$, $n = N/V$ is the number density of particles, and $\lambda = \sqrt{2\pi\hbar^2/m\tau}$ is the thermal de Broglie wavelength. The free energy $F = E - TS$ may be easily obtained by integrating the above P with respect to the volume V (since $P = -(\partial F/\partial V)_\tau$), and hence also the energy $E = -\tau^2[\partial(F/\tau)/\partial\tau]$. After subtracting out the energy of the perfect gas part (classical or quantal), one obtains the virial series for the interaction energy :

$$\frac{E_{int}}{V} = \frac{3}{2}n\tau \left[(n\lambda^3)\left\{a_2 - \frac{2}{3}\tau\frac{da_2}{d\tau}\right\} + (n\lambda^3)^2\left\{a_3 - \frac{2}{3}\tau\frac{da_3}{d\tau}\frac{1}{2}\right\} + \dots(n\lambda^3)^{(j-1)}\left\{a_j - \frac{2}{3}\tau\frac{da_j}{d\tau}\frac{1}{j-1}\right\} + \dots \right] \quad (2)$$

This expression is valid for a classical as well as a quantum gas, provided the appropriate classical or quantal virial coefficients are used. Whereas for a classical gas the virial coefficients may be expressed as integrals involving the interaction potential, for the quantum problem a solution of the j-body problem is needed to obtain a_j . Henceforth we shall denote the classical virial coefficients by $\mathcal{A}_j$ to differentiate from their quantum counterparts, to be still denoted as a_j . For an interesting example where classical and quantum results are very different, consider a hard sphere classical Boltzmann gas. For a hard-sphere diameter r_c , $\mathcal{A}_2 = \frac{2\pi}{3}(\frac{r_c}{\lambda})^3$, and the higher order $\mathcal{A}_j$'s may be expressed as $\mathcal{A}_j = \kappa(\mathcal{A}_2)^{j-1}$, where κ is a constant. Substituting this in Eq.(2), we see that the interaction energy of the classical hard sphere gas vanishes identically in each power of $(n\lambda^3)$. This, of course, is not the case for a bosonic or fermionic quantum gas at low temperatures.

We note that for a classical gas, $(n\lambda^3) \ll 1$, and the series (2) may be terminated after the first order term in most cases:

$$\frac{E_{int}}{V} = \frac{3}{2}n\tau \left[(n\lambda^3) \left\{ a_2 - \frac{2}{3}\tau \frac{da_2}{d\tau} \right\} \right] . \quad (3)$$

The classical second virial coefficient $\mathcal{A}_2$ is given by [4]

$$\mathcal{A}_2(\beta) = \frac{2\pi}{\lambda^3} \int_0^\infty dr \, r^2 (1 - \exp(-\beta V(r))) , \quad (4)$$

where $\beta = 1/\tau$, and $V(r)$ is the interatomic two-body potential. The quantum second virial coefficient requires a knowledge of the bound-state and the continuum spectra. For a one-component quantum gas, $b_2 = -a_2$ is given by [4]

$$b_2(\beta) = \sum_{l=0}^{\infty} (2l+1) b_2^{(l)} \quad (5)$$

$$b_2^{(l)}(\beta) = 2\sqrt{2} \left[\sum_n \exp(-E_{n,l}\beta) + \frac{1}{\pi} \int_0^\infty dk \, \frac{\partial \delta_l}{\partial k} \exp\left(-\frac{\hbar^2 k^2 \beta}{m}\right) \right] . \quad (6)$$

In the above, $E_{n,l}$'s are the two-body bound states (if any), and δ_l is the scattering phase shift in the l th partial wave. In the quantum case, for a Fermi gas of atoms interacting with a delta-function pseudo-potential, the spin-up atoms interact only with the spin-down ones, and one may consider a two-component (up, down) gas with $n_\uparrow = n_\downarrow = n/2$. In this situation, n^2 in (3) should be replaced by $n^2/4$. Often, the multiplicative prefactor in Eq.(6) is omitted, being transferred instead to the RHS of Eq.(3). The modified equation for a two-component quantum gas then reads

$$\frac{E_{int}}{V} = \frac{3}{2}n\tau \left[(n\lambda^3) \left\{ -\frac{b_2}{\sqrt{2}} + \frac{\sqrt{2}}{3}\tau \frac{db_2}{d\tau} \right\} \right] , \quad (7)$$

with

$$b_2^{(l)}(\beta) = \left[\sum_n \exp(-E_{n,l}\beta) + \frac{1}{\pi} \int_0^\infty dk \, \frac{\partial \delta_l}{\partial k} \exp\left(-\frac{\hbar^2 k^2 \beta}{m}\right) \right] . \quad (8)$$

Ho and Mueller [3] used the above formula for the quantum virial coefficient in the high temperature expression (7) to explain the low-temperature data. Two questions immediately arise regarding this treatment :

(a) For the quantum gas, since $(n\lambda^3)$ is larger than, or of order unity, higher order terms in Eq.(2) may be important. Presumably, near the Feshbach resonance, it is still the two-body term that dominates the interaction energy.

(b) One may also argue that while using the high-temperature expression (7) for the interaction energy, one should take the classical Eq.(4) for the second virial coefficient coefficient, instead of Eq.(8).

In the following, we attempt to answer the second question in some detail. Note that at low temperatures (large β), only small values of k contribute dominantly in the integrand in Eq. (6), and therefore only the $l = 0$ partial wave in the sum is important. Therefore, rather than the total classical $\mathcal{A}_2$ given by Eq. (4), we should consider its $l = 0$ classical component. Making the decomposition

$$\mathcal{A}_2(\beta) = \sum_{l=0}^{\infty} (2l+1) \mathcal{A}_2^{(l)} , \quad (9)$$

it may be verified easily (by replacing the sum over l by an integral), that

$$\mathcal{A}_2^{(l)} = \frac{1}{\lambda} \int_0^{\infty} dr \exp \left[-\frac{\hbar^2 l(l+1)}{mr^2} \beta \right] [1 - \exp(-\beta V(r))] . \quad (10)$$

One may enquire about the relation between the “semiclassical” expression (10) and the quantum counterpart $-b_2^{(l)}$. This question was actually answered nearly seventy years back by Kahn [5] in his doctoral dissertation. We shall give here a new compact derivation of his result, and for simplicity consider a potential with no bound states present. Using (8), and the relation $E = \frac{\hbar^2 k^2}{m}$, we have

$$b_2^{(l)}(\beta) = \left[\frac{1}{\pi} \int_0^{\infty} dk \frac{\partial \delta_l}{\partial k} \exp(-\frac{\hbar^2 k^2 \beta}{m}) \right] = \left[\frac{1}{\pi} \int_0^{\infty} dE \frac{\partial \delta_l}{\partial E} \exp(-\beta E) \right] . \quad (11)$$

With this definition of the virial coefficient, at a zero energy resonance, $b_2^{(0)} = 1/2$ or $-1/2$, according as the scattering length is negative (attractive interaction), or positive (repulsive interaction). We also note that $b_2^{(l)}(\beta)$ is just the Laplace transform of the derivative of the phase shift with respect to the energy. In the lowest order WKB approximation, the phase shift is given by [6]

$$[\delta_l(E)]_{WKB} = \sqrt{\frac{m}{\hbar^2}} \left[\left(\int_{r_l}^{\infty} \sqrt{E - V_l(r)} - \int_{r_0}^{\infty} \sqrt{E - \frac{\hbar^2 l(l+1)}{mr^2}} \right) dr \right] . \quad (12)$$

In the above, the effective potential $V_l(r)$ is defined as

$$V_l(r) = V(r) + \frac{\hbar^2 l(l+1)}{mr^2} . \quad (13)$$

In Eq.(12), r_l and r_0 are the classical turning points where the respective integrands go to zero. The derivative of the phase shift may be written as

$$\left[\frac{d\delta_l}{dE} \right]_{WKB} = \frac{1}{2} \sqrt{\frac{m}{\hbar^2}} \left[\left(\int_0^{\infty} \frac{\Theta(E - V_l(r))}{\sqrt{E - V_l(r)}} - \int_0^{\infty} \frac{\Theta(E - \frac{\hbar^2 l(l+1)}{2mr^2})}{\sqrt{E - \frac{\hbar^2 l(l+1)}{mr^2}}} \right) dr \right] \quad (14)$$

in terms of the unit step function $\Theta(x)$. By noting that the Laplace transform of $\frac{\Theta(E-\mu)}{\sqrt{E-\mu}}$ is $\sqrt{\frac{\pi}{\beta}} \exp(-\beta\mu)$, we immediately obtain the desired result

$$[b_2]_{WKB}^{(l)} = -\frac{1}{\sqrt{2}\lambda} \int_0^{\infty} dr \exp \left[-\beta \frac{\hbar^2 l(l+1)}{mr^2} \right] [1 - \exp(-\beta V(r))] = -\frac{\mathcal{A}_2^{(l)}}{\sqrt{2}} . \quad (15)$$

The multiplicative factor of $\frac{1}{\sqrt{2}}$ may be understood by noting that $b_2^{(l)}$ as defined by Eq.(8), is the quantum canonical two-body partition function for the *relative* motion in the l^{th} partial

wave, *without* the noninteracting continuum part. Noting that the radial motion in a given partial wave is one dimensional, the classical limit of $b_2^{(l)}$ is

$$Z_{cl}^{(l)} = \frac{1}{h} \int_{-\infty}^{\infty} dp_r \exp\left(-\frac{p_r^2 \beta}{m}\right) \int_0^{\infty} dr \left\{ \exp\left[-\beta\left(V(r) + \frac{\hbar^2 l(l+1)}{mr^2}\right)\right] - \exp\left[-\beta\left(\frac{\hbar^2 l(l+1)}{mr^2}\right)\right] \right\}, \quad (16)$$

$$= -\frac{1}{\sqrt{2\lambda}} \int_0^{\infty} dr \exp\left[-\beta\frac{\hbar^2 l(l+1)}{mr^2}\right] [1 - \exp(-\beta V(r))] = -\frac{\mathcal{A}_2^{(l)}}{\sqrt{2}}. \quad (17)$$

If the lowest order WKB approximation (14) for a given potential were exact, the classical and quantum (partial wave) virial coefficients would be related by $b_2^{(l)} = -\mathcal{A}_2^{(l)}/\sqrt{2}$. But in general, specially near the Feshbach resonance, the WKB approximation for the phase shift is not accurate. Moreover, the universality property of the quantum $b_2^{(0)} = \pm 1/2$ at the resonance is not reflected by its classical or WKB counterpart. The latter invariably depends on temperature as well as the parameters of the potential (even at resonance). As an illustration, consider an attractive square-well potential of depth $-V_0$, and range R for the two-body interaction. When the dimensionless “strength factor” $s = \frac{4}{\pi^2} \frac{mV_0}{\hbar^2} R^2$ is unity, there is an s-wave zero-energy bound state, with $b_2^{(0)} = \frac{1}{2}$. If, on the other hand, we were to use the WKB expression (15), we would obtain $[b_2^{(0)}]_{WKB} = \frac{R}{\sqrt{2\lambda}} (\exp(\beta V_0) - 1)$. Unlike the exact quantum result, this depends both on temperature and the potential parameters, so that universality is lost.

The only exception to this rule is for a scale-independent inverse square potential of the type

$$V(r) = \frac{\hbar^2 \alpha^2}{m r^2}, \quad (18)$$

where α is a dimensionless (real) parameter. It is known that the WKB method gives the exact phase shift [7] for such a potential, and it is easy to check that the classical virial coefficient, evaluated using Eq.(10), is

$$(\mathcal{A}_2^{(0)})_{res} = \frac{|\alpha|}{\sqrt{2}}. \quad (19)$$

Using Eq.(17) we see that for $\alpha^2 = 1$, we get $b_2^{(0)} = -1/2$, as is appropriate for a repulsive potential at a zero-energy resonance.

We now show that the classical form of the second virial coefficient may also be recovered from the quantum expression through an exact mapping near the zero-energy resonance. To this end, we start with Eq.(11) for the $l = 0$ partial wave,

$$b_2^{(0)}(\beta) = \left[\frac{1}{\pi} \int_0^{\infty} dk \frac{d\delta_0}{dk} \exp\left(-\frac{\hbar^2 k^2 \beta}{m}\right) \right]. \quad (20)$$

Near zero-energy ($k \rightarrow 0$), we may use the shape-independent effective range expansion, and for $l = 0$ it is given by

$$k \cot \delta_0 = -\frac{1}{a} + \frac{1}{2} r_0^2 + \dots \quad (21)$$

In the above, a is the scattering length that approaches $\pm\infty$ near a resonance, and r_0 is the effective range. The virial coefficient near the resonance is governed by a alone, and we may ignore the second term. In that case, using (20), we get

$$b_2^0 = -\frac{1}{\pi} \int_0^\infty dk \frac{a}{1+a^2k^2} \exp\left(-\frac{\hbar^2 k^2 \beta}{m}\right). \quad (22)$$

This may be rewritten as

$$b_2^0 = \frac{1}{\pi} \int_0^\infty dk \frac{a}{1+a^2k^2} \left(1 - \exp\left(-\frac{\hbar^2 k^2 \beta}{m}\right)\right) - \frac{1}{\pi} \int_0^\infty dk \frac{a}{1+a^2k^2}. \quad (23)$$

The second term yields $-sgn(a) \frac{1}{2}$. As $a \rightarrow \pm\infty$, we may then write

$$b_2^{(0)} \simeq \frac{sgn(a)}{|a|\pi} \int_0^\infty \frac{dk}{k^2} \left(1 - \exp\left(-\frac{\hbar^2 k^2 \beta}{m}\right)\right) - sgn(a) \frac{1}{2} \quad (24)$$

By making the transformation $k = \alpha/x$, we get

$$b_2^{(0)} \simeq \frac{sgn(a)}{|a|\pi\alpha} \int_0^\infty \left(1 - \exp\left(-\frac{\hbar^2 \alpha^2 \beta}{mx^2}\right)\right) dx - sgn(a) \frac{1}{2} \quad (25)$$

We have already noted, from Eq.(19), that the second term on the RHS may be reproduced classically by an inverse square potential with $\alpha = 1$ for the repulsive case. The first term on the RHS, on performing the integral, is independent of α , and gives a correction of order $\hbar$ to the classical value in the neighbourhood of the resonance. We may thus write the above equation in the form

$$b_2^{(0)} \simeq sgn(a) \left(-\frac{\mathcal{A}_2^{(0)}}{\sqrt{2}}\right)_{res} \left[1 - \frac{\sqrt{2}\lambda}{|a|\pi}\right], \quad (26)$$

where $\mathcal{A}_2^{(0)}$ is obtained from Eq.(10) with the inverse square potential (18) of strength $\alpha = 1$.

We have shown above that as the Feshbach resonance is approached from the repulsive side, the two-body physics may be simulated by a universal inverse square potential of unit strength. This does not appear to hold on the attractive side. The inverse square potential in one space dimension is known as the Calogero-Sutherland potential [8], and has been thoroughly studied, including its two-point correlation function [9]. The integral of the latter is related to the isothermal compressibility, and can be computed numerically [10]. Its experimental measurement near the Feshbach resonance may be of interest, specially in view of the findings in this paper. The other intriguing point is the so-called Efimov effect [11]. When the two-body scattering length approaches infinity from the *attractive side*, a very large number of three-body bound states appear. More over, Efimov has shown that the universal effective three-body interaction in this case is $-\frac{s_0^2 \hbar^2}{2mR^2}$, where $R^2 = 2(r_{12}^2 + r_{23}^2 + r_{31}^2)/3$, and $s_0^2 \simeq 1$. We find, from the two-body analysis, that an inverse square two-body effective potential also arises at Feshbach resonance, albeit on the repulsive side. More research in this direction is planned.

We thank Don Sprung for going through the manuscript carefully. One of the authors (R.K.B.) acknowledges NSERC (Canada) for financial support.

REFERENCES

- [1] S. Inouye *et al.*, Nature (London) **392**, 151 (1998); Ph. Courteille *et al.*, Phys. Rev. Lett. **81**, 69 (1998).
- [2] G.A. Baker, Phys. Rev. C**60**, 054311 (1999); H. Heiselberg, Phys. Rev A**63**, 043606 (2001); T-L Ho, Phys. Rev. Lett. **92**, 090402 (2004).
- [3] T-L Ho and E.J. Mueller Phys. Rev. Lett. **92**, 160404 (2004).
- [4] R.K. Pathria, *Statistical Mechanics*, (Pergamon Press, Oxford, 1972) chapter 9.
- [5] Boris Kahn, Doctoral Dissertation, University of Utrecht, 1938 (reproduced in Studies in Statistical Mechanics, Vol.3, eds. J. De Boer and G.E. Uhlenbeck, North Holland, Amsterdam, 1965) p. 281.
- [6] A. Messiah, *Quantum Mechanics, Vol.1* (North Holland, Amsterdam, 1961) p.410.
- [7] H. Guérin, J. Phys. B**29**, 1285 (1996).
- [8] F. Calogero, J. Math. Phys. **10**, 2191 (1969); *ibid* **10**, 2197 (1969); B. Sutherland, J. Math. Phys. **12**, 246 (1971).
- [9] V.Ya. Krivnov and A.A. Ovchinnikov, JETP **55** 162 (1982); D. Sen and R.K. Bhaduri, Can. J. Phys. **77**, 327 (1999).
- [10] R.K. Bhaduri, M.V.N. Murthy, and M.N. Tran, J. Phys. B**35**, 2817 (2002).
- [11] V. Efimov, Sov. J. Nuclear Phys. **12** 589 (1971); Phys. Lett. B**33**, 563 (1973).